

Santos, X. 2008. *Natrix maura* (Linnaeus, 1758). 176-177. In: Loureiro, A., Ferrand de Almeida, N., Carretero, M. A. & Paulo, O.S. (eds.), *Atlas dos Anfíbios e Répteis de Portugal*. Instituto da Conservação da Natureza e da Biodiversidade. Lisboa.

Santos, X. 2014. *Natrix maura* (Linnaeus, 1758). 820-842. In:

Salvador, A. (coordinador), *Reptiles, 2ª edición revisada y aumentada*. Ramos, M.A. et al. (eds.), *Fauna Ibérica, vol. 10*. Consejo Superior de Investigaciones Científicas. Madrid.

Schleich, H., Kästle, W. & Kabisch, K. 1996. *Amphibians and Reptiles of North Africa*. Koeltz Sci. Books. Koenigstein.

## The yellow-legged gull, *Larus michahellis*, an occasional predator of *Timon lepidus* in the Atlantic Islands National Park (Galicia, NW Spain)

Vicente Piorno<sup>1\*</sup>, José-Antonio Fernández<sup>1</sup> & Guillermo Velo-Antón<sup>2</sup>

<sup>1</sup> Parque Nacional de las Islas Atlánticas de Galicia. Consellería de Medio Ambiente e Ordenación do Territorio, Xunta de Galicia. R. Oliva, 3. 36202 Vigo. Spain. C.e.: vicente.piorno.gonzalez@xunta.es

<sup>2</sup> CIBIO/InBIO, Centro de Investigación em Biodiversidade e Recursos Genéticos da Universidade do Porto. Instituto de Ciências Agrárias de Vairão. R. Padre Armando Quintas. 4485-661 Vairão. Portugal.

**Fecha de aceptación:** 3 de noviembre de 2015.

**Key words:** islands, *Larus michahellis*, predation, *Timon lepidus*.

**RESUMEN:** El lagarto ocelado (*Timon lepidus*) mantiene poblaciones de gran interés en varias islas, tanto por su estatus de conservación como por las particulares interacciones ecológicas que en ellas se desarrollan. En el caso del Parque Nacional de las Islas Atlánticas de Galicia, esta especie convive con una importante colonia de cría de gaviota patiamarilla (*Larus michahellis*). Desde el año 2009 hasta 2015 hemos detectado en este Parque Nacional siete casos de consumo de *T. lepidus* por *L. michahellis*. Se trata de una conducta nunca antes documentada y de reciente aparición, que podría estar relacionada con los efectos que el cierre de vertederos y la reducción de los descartes pesqueros están teniendo en esta especie de gaviota. El carácter ocasional de esta depredación no hace esperar efectos apreciables en estas poblaciones insulares de lagarto ocelado.

The ocellated lizard, *Timon lepidus* (Daudin, 1802), is a Mediterranean reptile widespread throughout most of the Iberian Peninsula, and patchily distributed in southern France and northwestern Italy (Sillero et al., 2014). It occurs in a wide variety of open habitats and along a broad altitudinal range. Due to its basking behaviour and relatively large size, it is part of the diet of a high number of medium and large sized Iberian vertebrates. These predators are well documented, and include mostly birds. A literature review of avian predation on this species (Martín & López, 1996) shows the Falconiformes as com-

mon predators of *T. lepidus*, while the orders Strigiformes, Ciconiiformes and Charadriiformes are occasional predators. Only one species, *Gelochelidon nilotica*, has been reported as a predator of *T. lepidus* within Charadriiformes (Martín & López, 1990).

Throughout its range, *T. lepidus* occurs on some Atlantic and Mediterranean islands, some of which had experienced marked population declines (Mateo, 2015). Island native populations are usually of high conservation interest due to the combination of singular characteristics of islands ecosystems (e.g., isolation, age, size), with particular species

interactions that are absent or less common in mainland populations (e.g., due to differences in trophic resources and presence of predators) (Wardle, 2002). For example, some island populations of *T. lepidus* show differences in antipredator behaviour (Vicente, 1989), social behaviour (Galán, 2000) and levels of frugivory (Piazzon *et al.*, 2012), when compared to their mainland counterparts.

The Atlantic Islands National Park (Galicia, NW Spain) represents a relevant example of these island populations of *T. lepidus*. This species is present on the main islands of the four archipelagos of the Park (Galán, 2003; Figure 1), and a subspecies endemic to one of them (Sálvora), *T. lepidus oteroi*, has been described (Castroviejo & Mateo, 1998). Two other archipelagos, Cíes and Ons, harbour populations with very similar morphological characteristics, probably belonging to the same subspecies (Galán, 2003), although further genetic and morphological studies are needed to support the validity of this subspecies across all islands. These populations have been listed as Vulnerable (Xunta de Galicia, 2007) due to their unique taxonomic status and the decline experienced in the past on some of the islands.

The Atlantic Islands National Park also harbours important breeding colonies of Yellow-legged gull (*Larus michahellis*), which are among the largest in the world (Molina & Bermejo, 2009). The coexistence of *T. lepidus* and *L. michahellis* in such a relatively small territory and under the particular conditions of an island ecosystem provides the opportunity for interactions that are less likely on the mainland.

After receiving some occasional reports of gulls feeding on lizards from several islands of the Park, we decided to interview all the field personnel of the National Park. There are a total of 26 workers working in shifts, ensuring a permanent presence in all of the archipelagos since the protection of these islands (1980 in the case of Cíes, 1982 in the case of Ons, and 2002 in the case of Sálvora and Cortegada). We asked all of the workers if they had observed this behaviour, and, when the answer was affirmative, if they had observed the lizard previously alive. We also asked for available photographs of these interactions, and the exact place and date of each event. We checked all the reported events with the observers to discard possible duplicated observations.

We registered seven events of *L. michahellis* feeding on *T. lepidus*. In four of these cases, the



**Figure 1:** Distribution of *T. lepidus* (Galán, 2007) and *L. michahellis* (Pérez *et al.*, 2012) in the Atlantic Islands National Park.

**Figura 1:** Distribución de *T. lepidus* (Galán, 2007) y *L. michahellis* (Pérez *et al.*, 2012) en el Parque Nacional de las Islas Atlánticas.

observer could confirm that the lizard was alive before the event. We, however, cannot completely rule out some possible cases of scavenging in the other three, considering that the gulls were already feeding on a dead lizard when this behaviour was observed. The first of these events was registered in 2009 and the last one in 2015. We obtained photographs and dates for three of these cases (Figure 2), all of them happening in the second half of May. For the other four cases, we only obtained the year. There is no appreciable annual trend in the number of observed events (Figure 3). All the cases were observed in Cíes and Sálvora, the two archipelagos of the National Park with the highest abundances of the two species (Table 1). The number of events was highest in Cíes, where there are more potential observers (National Park staff). There are neither recorded cases in Ons, where gulls and lizards are less abundant, nor in Cortegada, where there are no breeding gulls.

There are reports of predation of *L. michabellis* on other small lizards (Martín & López, 1990; Castilla & Labra, 1998; Vervust *et al.*, 2007; Pérez-Mellado *et al.*, 2014). It has also been suggested, without specific evidence, that gulls can predate on *T. lepidus* (Bischoff *et al.*, 1984; Cheylan & Grillet, 2004; Doré *et al.*, 2015). However, to our knowledge, this is the first confirmed case of *L. michabellis* preying on a lizard as large as *T. lepidus*.

We believe that this behaviour is not very common on these islands, taking into account that, since the first cases, the National Park workers have been very attentive to these events. In addition, we have been carrying out analysis of gull pellets over several years, without having found lizard remains in them (Atlantic Islands National Park, unpublished data). Considering the high number of gulls on the islands and the low number of events observed, they could represent a case of specia-




**Figure 2:** *Larus michabellis* feeding on *Timon lepidus*. (a) Sálvora Island, 31/5/2009; (b) Faro Island, Cíes, 13/5/2013; (c) Monteagudo Island, Cíes, 30/5/2015.

**Figura 2:** *Larus michabellis* alimentándose de *Timon lepidus*. (a) Sálvora Island, 31/5/2009; (b) Faro Island, Cíes, 13/5/2013; (c) Monteagudo Island, Cíes, 30/5/2015.

lized predation of a few individuals, a common behaviour in this species (Ceia *et al.*, 2014).

All the events with a known date occurred in May, when these lizards show a seasonal peak in their activity (Mateo, 2015). Therefo-



**Figure 3:** Annual distribution of the observed cases of *Larus michahellis* feeding on *T. lepidus*.

**Figura 3:** Distribución anual de los casos observados de *Larus michahellis* alimentándose de *T. lepidus*.

re, this behaviour has probably been facilitated in the National Park by the coexistence in time and space of these two species, when both gulls and lizards breed during spring, and the former actively search for food to feed their chicks. Population density seems also to have an effect, considering that observed interactions occurred in the islands where both species are most abundant.

It is also worth noting the recent emergence of this species interaction. There has been a continued presence of conservationists and researchers visiting these islands since their protection, with regular visits by ornithologists and herpetologists that have not previously reported cases of such interaction. One possible explanation is the recent decrease in food resources available to *L. michahellis*. This is an opportunistic species that frequently uses two anthropogenic food sources, garbage dumps and fishing discards, which, due to its easy access, led to a marked increase in its populations at the end of the XX century (Blokpoel & Spaans, 1991, and references therein). Recently, two European Union environmental action plans aiming at reducing fishing discards and closing garbage dumps have decreased the availability of these resources, with a considerable impact on gulls' diet (Ramos *et al.*, 2009). The population of *L. michahellis* in the

Park seems to be suffering the consequences of this reduction in food availability. The number of breeding pairs has experienced a decrease of 65.8% over the last 10 years (Atlantic Islands National Park, unpublished data). This situation has led to changes in the feeding ecology of *L. michahellis*, increasing the diversity of its diet (Duhem *et al.*, 2003) and the levels of predation on smaller species (Oro & Martínez-Abraín 2007). This idea is reinforced by at least two other recent documented cases of *L. michahellis* preying on unusual preys (Báez, 2015; Ros, 2015).

Overall, these observations point to *L. michahellis* as an occasional predator of *T. lepidus*, with an unlikely role in the population decrease of *T. lepidus* registered in some of these islands. *Larus michahellis* has been unfairly blamed in the past for negatively affecting other species (Oro & Martínez-Abraín, 2007; Pérez-Mellado *et al.*, 2014). However, the real threats to *T. lepidus* in the Park have already been identified. Introduced domestic cats, *Felis silvestris catus*, considered the most successful and damaging invader on islands (Bonnaud *et al.*, 2011), are the main cause of concern (Galán, 2003, 2007). Moreover, introduced American mink, *Neovison vison*, and black rat, *Rattus rattus*, might be potential threats for amphibians and reptile populations of these islands (Velo-Antón & Cordero-Rivera, 2011; Romero, 2015). Eradication campaigns of cats

**Table 1:** Observed events, number of observers and abundance of *T. lepidus* (Galán, 2007) and *L. michabellis* (Pérez et al., 2012) on the islands of the Atlantic Islands National Park.**Tabla 1:** Eventos observados, número de observadores y abundancia de *T. lepidus* (Galán, 2007) y *L. michabellis* (Pérez et al., 2012) en las islas del Parque Nacional de las Islas Atlánticas.

	Observed events	Permanent staff	<i>T. lepidus</i> abundance		<i>L. michabellis</i> abundance	
			Individuals/ha	SE	Breeding pairs	Density (pairs/ha)
Cíes - Monteagudo	1	4	58,7	22,1	3.771	19,95
Cíes - Faro	5	4	96,4	27,0	1.271	12,67
Cíes - San Martiño	0	0	35,0	10,0	2.423	15,36
Ons	0	4	15,5	3,9	2.246	5,27
Sálvora	1	1	57,0	26,7	4.786	19,99
Cortegada	0	1	NA	NA	0	0,00

and minks have been successfully conducted since 2009 by the staff of the National Park and they are not currently considered a major threat. Both *L. michabellis* and *T. lepidus* have been inhabiting these islands for a long time, and there is no evidence to think that this coexistence would threaten *T. lepidus* populations in the near future.

**ACKNOWLEDGMENTS:** We thank all the interviewed employees of the National Park for their help in this study, and in particular, J. Bernárdez, M. Caneda, M. Chamorro, P. Domínguez, J.M. López, J. Ledo and P. Vázquez for their observations of this predation behaviour. M. Chamorro, P. Domínguez and P. Vázquez kindly allowed us to publish their photographs.

## REFERENCES

- Báez, J.C. 2015. Depredación sobre el galápagos invasor *Trachemys scripta* por *Larus cachinnans*. *Boletín de la Asociación Herpetológica Española*, 26: in press.
- Bischoff, W., Cheylan, M., & Böhme, W. 1984. *Lacerta lepida* Daudin, 1802 - Perleidechse. 181-210. In: Böhme, W. (ed.), *Handbuch der Reptilien und Amphibien Europas. Band 2/I: Echsen (Sauria) II (Lacertidae II: Lacerta)*. Aula-Verlag, Wiesbaden.
- Blokpoel, H. & Spaans, A. 1991. Introductory remarks: superabundance in gulls: causes, problems and solutions. *Acta Congressus Internationalis Ornithologici*, 20: 2361–2364.
- Bonnaud, E., Medina, F.M., Vidal, E., Nogales, M., Thershy, B., Zavaleta, E., Donlan, C.J., Keitt, B., Le Corre, M. & Horwath, S.W. 2011. The diet of feral cats on islands: a review and a call for more studies. *Biological Invasions*, 13: 581–603.
- Castilla, A.M. & Labra, A. 1998. Predation and spatial distribution of the lizard *Podarcis hispanica atrata*: an experimental approach. *Acta Oecologica*, 19: 107–114.
- Castroviejo, J. & Mateo, J.A. 1998. Una nueva subespecie de *Lacerta lepida* Daudin, 1802 (Sauria, Lacertidae) para la isla de Sálvora (España). *Publicaciones de la Asociación de Amigos de Doñana*, 12: 1–21.
- Ceja, F.R., Paiva, V.H., Fidalgo, V., Morais, L., Baeta, A., Crisóstomo, P., Mourato, E., Garthe, S., Marques, J.C. & Ramos J.A. 2014. Annual and seasonal consistency in the feeding ecology of an opportunistic species, the yellow-legged gull *Larus michabellis*. *Marine Ecology Progress Series*, 497: 273–284.
- Cheylan, M. & Grillet, P. 2004. *Le Lézard ocellé*. Éditions Belin Éveil Nature. Tours, France.
- Doré, F., Cheylan, M., Grillet, P., Dusoulier, F. & Corti, C. 2015. *Le lézard ocellé : Un géant sur le continent européen*. Biotope Editions. Mèze, France.
- Duhem, C., Vidal, E., Roche, P. & Legrand, J. 2003. Island breeding and continental feeding: How are diet patterns in adult yellow-legged gulls influenced by landfill accessibility and breeding stages? *Ecoscience*, 10: 502–508.
- Galán, P. 2000. Conducta gregaria del lagarto ocelado, *Lacerta lepida*, en las islas Cíes (Galicia). *Boletín de la Asociación Herpetológica Española*, 11: 80–82.
- Galán, P. 2003. *Anfibios y reptiles del Parque Nacional de las Islas Atlánticas de Galicia*. Ministerio de Medio Ambiente. Madrid.
- Galán, P. 2007. *Plan de manejo y gestión de los reptiles del Parque Nacional Marítimo Terrestre de las Islas Atlánticas de Galicia*. Unpublished report. Atlantic Islands National Park. Vigo, Spain.
- Martín, J. & López, P. 1990. Amphibians and reptiles as prey of birds in southwestern Europe. *Smithsonian Herpetological Information Service*, 82: 1–43.
- Martín, J. & López, P. 1996. Avian predation on a large lizard (*Lacerta lepida*) found at low population densities in Mediterranean habitats: An analysis of bird diets. *Copeia*, 1996: 722–726.
- Mateo, J.A. 2015. Lagarto ocelado - (*Timon lepidus*). In: Salvador, A. & Marco, A. (eds.), *Enciclopedia virtual de los vertebrados españoles*. Museo Nacional de Ciencias Naturales. Madrid. <http://www.vertebradosibericos.org/reptiles/iden-

- tificacion/timlepid.html> [Accessed: 10 October 2015].
- Molina, B. & Bermejo, A. 2009. La gaviota patiamarilla. 50–111. In: Molina, B. (ed.), *Gaviotas reidora, sombría y patiamarilla en España. Población en 2007-2009 y método de censo*. SEO/BirdLife. Madrid.
- Oro, D. & Martínez-Abraín, A. 2007. Deconstructing myths on large gulls and their impact on threatened sympatric waterbirds. *Animal Conservation*, 10: 117–126.
- Pérez, C., Barros, A., Velando, A., & Munilla, I. 2012. *Seguimento das poboacións reproductoras de corvo mariño (Phalacrocorax aristotelis) e gaviota patimarela (Larus michahellis) do Parque Nacional das Illas Atlánticas de Galicia*. Unpublished report. Atlantic Islands National Park. Vigo, Spain.
- Pérez-Mellado, V., Garrido, M., Ortega, Z., Pérez-Cembranos, A., & Mencía, A. 2014. The yellow-legged gull as a predator of lizards in Balearic Islands. *Amphibia-Reptilia*, 35: 207–213.
- Piazzon, M., Larrinaga, A.R., Rodríguez-Pérez, J., Latorre, L., Navarro, L., & Santamaría, L. 2012. Seed dispersal by lizards on a continental-shelf island: predicting interspecific variation in seed rain based on plant distribution and lizard movement patterns: Seed dispersal by lizards. *Journal of Biogeography*, 39: 1984–1995.
- Ramos, R., Ramírez, F., Sanpera, C., Jover, L. & Ruiz, X. 2009. Diet of Yellow-legged Gull (*Larus michahellis*) chicks along the Spanish Western Mediterranean coast: the relevance of refuse dumps. *Journal of Ornithology*, 150: 265–272.
- Romero, R. 2015. *Actuaciones de control y seguimiento de Visón americano y Gato asilvestrado en el parque Nacional de las Islas Atlánticas de Galicia*. Unpublished report. Atlantic Islands National Park. Vigo, Spain.
- Ros, J.A. 2015. Depredación de gaviota patiamarilla (*Larus michahellis*) sobre *Natrix maura* en Cartagena (Murcia). *Boletín de la Asociación Herpetológica Española*, 26: 32–33.
- Sillero, N., Campos, J., Bonardi, A., Corti, C., Creemers, R., Crochet, P.-A., Isailović, J.C., Denoël, M., Ficetola, G.F., Gonçalves, J., Kuzmin, S., Lymberakis, P., Pous, P. de, Rodríguez, A., Sindaco, R., Speybroeck, J., Toxopeus, B., Vieites, D.R. & Vences, M. 2014. Updated distribution and biogeography of amphibians and reptiles of Europe. *Amphibia-Reptilia*, 35: 1–31
- Velo-Antón, G. & Cordero-Rivera, A. 2011. Predation by invasive mammals on an insular viviparous population of *Salamandra salamandra*. *Herpetology Notes*, 4: 299–301.
- Vervust, B., Grbac, I. & Van Damme, R. 2007. Differences in morphology, performance and behaviour between recently diverged populations of *Podarcis sicula* mirror differences in predation pressure. *Oikos*, 116: 1343–1352.
- Vicente, L. 1989. Novos dados sobre o comportamento de *Lacerta lepida* (Daudin, 1802) (Sauria-Lacertidae). Uma população insular Ilha da Berlenga (Portugal). *Treballs de la Societat Catalana d'Ictiologia i Herpetologia*, 2: 232–249.
- Wardle, D. A. 2002. Islands as model systems for understanding how species affect ecosystem properties. *Journal of Biogeography*, 29: 583–591.
- Xunta de Galicia. 2007. Decreto 88/2007 do 19 de abril, polo que se regula o Catálogo Galego de Especies Amenazadas. *Diario Oficial de Galicia*, 89: 7409–7423.

## *Pseudocerastes urarachnoides*: the ambush specialist

Gabriel Martínez del Marmol<sup>1</sup>, Omid Mozaffari<sup>2</sup> & Javier Gállego<sup>3</sup>

<sup>1</sup> Cl. Pedro Antonio de Alarcon, 34. 18002 Granada. Spain. C.e.: gabrimtnez@gmail.com

<sup>2</sup> Apt. 1, No. 34. Darougar cplx., 1st Milad. Eram Blvd. Mehrshahr, Alborz. Iran.

<sup>3</sup> Carretera Motores, 211. 04720 Aguadulce. Almeria. Spain.

**Fecha de aceptación:** 3 de noviembre de 2015.

**Key words:** *Pseudocerastes*, tail ornament, behavior, caudal luring, Viperidae.

**RESUMEN:** *Pseudocerastes urarachnoides* es una especie recientemente descrita en Oriente Medio, y los datos sobre su ecología y distribución todavía son escasos. Durante un viaje herpetológico a Irán en la primavera de 2014 pudieron observarse dos especímenes de esta especie en su hábitat natural. En este artículo se presentan datos sobre los hábitos de caza y hábitats usados, información importante para la conservación y el manejo de la especie.

Vipers of the genus *Pseudocerastes*, Boulenger 1896 are sometimes known as “false horned vipers” due to the presence of horn-like projections on both sides above the eye. These projections are formed by several small imbricate scales

and their tips end in two tiny scales. In contrast, the “true horned vipers” *Cerastes cerastes* (Linnaeus, 1758) and *Cerastes gasperettii* Leviton & Anderson, 1967 have supraorbital “horns”, each one consisting in a single elongated scale. Three